

ACEPROMAZINE 1 mg/mL		Dosage (IM or SQ) 0.05 mg/kg, capped at 1mg at 31≥ lbs (14≥ kg)	
Weight (Kgs)	mL	Weight (Lbs)	
1.0 - 2.5	0.1	2.2 - 5.5	
3.0 - 4.5	0.2	6.6 - 10.0	
5.0 - 9.0	0.5	11.0 - 19.8	
9.5 - 13.5	0.7	21.0 - 29.7	

ACEPROMAZINE 10 mg/mL		Dosage (IM or SQ) 0.05 mg/kg, capped at 1mg at 31≥ lbs (14≥ kg)	
Weight (Kgs)	mL	Weight (Lbs)	
14.0 ≥	0.10	30.8 ≥	

HYDROMORPHONE 2 mg/mL		Dosage (IM or SQ) 0.1 mg/kg	
Weight (Kgs)	mL	Weight (Lbs)	
1 - 2.5	0.1	2.0 - 5.5	
3 - 4.5	0.2	6.0 - 9.9	
5 - 7	0.3	11.0 - 15.0	
7.5 - 9	0.4	16.5 - 20.0	
9.5 - 11.5	0.5	20.9 - 25.3	
12 - 13.5	0.65	26.4 - 29.7	
14 - 16	0.75	30.8 - 35.0	
16.5 - 18	0.85	36.3 - 40.0	
18.5 - 20.5	1.0	40.7 - 45.1	
21 - 23	1.1	46.0 - 50.0	
23.5 - 25	1.2	51.7 - 55.0	
25.5 - 27	1.3	56.1 - 60.0	
27.5 - 29.5	1.4	60.5 - 64.9	
30 - 32	1.5	66.0 - 70.4	
32.5 - 34	1.6	71.5 - 75.0	
34.5 - 36.5	1.7	75.9 - 80.3	
37 - 38.5	1.9	81.0 - 84.7	
39 - 41	2.0	86.0 - 90.0	
41.5 - 43	2.1	91.3 - 95.0	
43.5 - 45.5	2.2	95.7 - 100.1	
46 - 48	2.3	101.0 - 105.0	
48.5 - 50	2.4	106.7 - 110.0	
50.5 - 55	2.6	111.1 - 120.0	

Consider giving premed intramuscularly (IM) vs. subcutaneously (SQ) to increase intensity & onset of analgesia & sedation.