

Alternative premedication if morphine is unavailable

ACEPROMAZINE 1 mg/mL		Dosage (IM or SQ) 0.05 mg/kg, capped at 1mg at 31≥ lbs (14≥ kg)	
Weight (Lbs)		mL	Weight (Kgs)
2 - 5		0.1	0.9 - 2.3
6 - 10		0.2	2.7 - 4.5
11 - 20		0.5	5.0 - 9.1
21 - 30		0.7	9.5 - 13.6

ACEPROMAZINE 10 mg/mL		Dosage (IM or SQ) 0.05 mg/kg, capped at 1mg at 31≥ lbs (14≥ kg)	
Weight (Lbs)		mL	Weight (Kgs)
31 ≥		0.10	14.1 ≥

HYDROMORPHONE 2 mg/mL		Dosage (IM or SQ) 0.1 mg/kg	
Weight (Lbs)		mL	Weight (Kgs)
2 - 5		0.1	1 - 2
6 - 10		0.2	3 - 5
11 - 15		0.3	5 - 7
16 - 20		0.4	7 - 9
21 - 25		0.5	10 - 11
26 - 30		0.65	12 - 14
31 - 35		0.75	14 - 16
36 - 40		0.85	16 - 18
41 - 45		1.0	19 - 20
46 - 50		1.1	21 - 23
51 - 55		1.2	23 - 25
56 - 60		1.3	25 - 27
61 - 65		1.4	28 - 30
66 - 70		1.5	30 - 32
71 - 75		1.6	32 - 34
76 - 80		1.7	35 - 36
81 - 85		1.9	37 - 39
86 - 90		2.0	39 - 41
91 - 95		2.1	41 - 43
96 - 100		2.2	44 - 45
101 - 105		2.3	46 - 48
106 - 110		2.4	48 - 50
111 - 120		2.6	50 - 55

Consider giving premed intramuscularly (IM) vs. subcutaneously (SQ) to increase intensity and onset of analgesia and sedation.