

For IT use, double the volume with saline			
EPINEPHRINE	Dosage		
	0.01 mg/kg (IV)	0.05 mg/kg (IT)	Weight (Lbs)
Weight (Kgs)	Concentration = 1:1000	Concentration = 1:1000	
1.0 - 2.5	0.03	0.15	2 - 5
3.0 - 4.5	0.04	0.2	6 - 10
5.0 - 9.5	0.05	0.25	11 - 21
10.0 - 14.5	0.10	0.5	22 - 32
15.0 - 19.5	0.15	0.75	33 - 43
20.0 - 24.5	0.20	1.0	44 - 54
25.0 - 29.5	0.25	1.25	55 - 65
30.0 - 34.5	0.30	1.5	66 - 76
35.0 - 39.5	0.35	1.75	77 - 87
40.0 - 44.5	0.40	2.0	88 - 98

For prolonged recovery, extreme sedation, emergencies, etc.			
NALOXONE 0.4 mg/mL	Dosage (IV, IM, or SQ)		
	0.01 mg/kg	0.04 mg/kg	Weight (Lbs)
Weight (Kgs)	mL	mL	
0.9	0.02	0.09	2
1.4	0.03	0.14	3
1.8	0.05	0.18	4
2.3	0.06	0.23	5
2.7	0.07	0.27	6
3.2	0.08	0.32	7
3.6	0.09	0.36	8
4.1	0.10	0.41	9
4.5	0.11	0.45	10
5 - 7	0.13	0.50	11 - 15
7 - 10	0.18	0.73	16 - 20
10 - 12	0.24	0.95	21 - 25
12 - 14	0.30	1.18	26 - 30
14 - 16	0.35	1.41	31 - 35
16 - 18	0.41	1.64	36 - 40
18 - 20	0.45	1.80	41 - 45
20 - 23	0.50	2.00	46 - 50
23 - 25	0.58	2.32	51 - 55
25 - 27	0.64	2.55	56 - 60
27 - 30	0.68	2.70	61 - 65
30 - 32	0.75	3.00	66 - 70
32 - 34	0.81	3.23	71 - 75
34 - 36	0.85	3.40	76 - 80
36 - 39	0.90	3.60	81 - 85
39 - 41	0.98	3.91	86 - 90
41 - 45	1.03	4.14	91 - 100
45 - 50	1.13	4.50	101 - 110
50 - 55	1.26	5.05	111 - 120

Atropine & Antisedan Charts on Reverse

For IT use, double the volume with saline			
ATROPINE 0.5 mg/mL	Dosage		
	0.04 mg/kg (IV)	0.2 mg/kg (IT)	
Weight (Kgs)	mL	mL	Weight (Lbs)
2.0 - 4.5	0.2	1.0	2 - 10
5.0 - 9.5	0.4	2.0	11 - 21
10.0 - 14.5	0.8	4.0	22 - 32
15.0 - 19.5	1.2	6.0	33 - 43
20.0 - 24.5	1.6	8.0	44 - 54
25.0 - 29.5	2.0	10.0	55 - 65
30.0 - 34.5	2.4	12.0	66 - 76
35.0 - 39.5	2.8	14.0	77 - 87
40.0 - 44.5	3.2	16.0	88 - 98

ANTISEDAN 5 mg/mL	Dosage (IM) 0.1 mg/kg	
	For Dexdomitor reversal	
Add 0.2 ml of saline to each dose of Antisedan for feline reversal		
Weight (Kgs)	mL	Weight (Lbs)
0.9 - 1.1	0.02	2 - 2.4
1.1 - 1.3	0.02	2.5 - 2.9
1.4 - 1.5	0.03	3 - 3.4
1.6 - 1.8	0.03	3.5 - 3.9
1.8 - 2.0	0.05	4 - 4.4
2.0 - 2.2	0.05	4.5 - 4.9
2.3 - 2.5	0.05	5 - 5.4
2.5 - 2.7	0.06	5.5 - 5.9
2.7 - 2.9	0.06	6 - 6.4
3.0 - 3.1	0.06	6.5 - 6.9
3.2 - 3.4	0.08	7 - 7.4
3.4 - 3.6	0.08	7.5 - 7.9
3.6 - 3.8	0.08	8 - 8.4
3.9 - 4.0	0.08	8.5 - 8.9
4.1 - 4.3	0.10	9 - 9.4
4.3 - 4.5	0.10	9.5 - 9.9
4.5 - 4.7	0.10	10 - 10.4
4.8 - 5.0	0.10	10.5 - 10.9
5.0 - 5.2	0.13	11 - 11.4
5.2 - 5.4	0.13	11.5 - 11.9
5.5 - 5.6	0.13	12 - 12.4
5.7 - 5.9	0.13	12.5 - 12.9
5.9 - 6.1	0.13	13 - 13.4
6.1 - 6.3	0.15	13.5 - 13.9

Antisedan may be administered IM.
The patient must be warm and hydrated.
Adding 0.2 mL saline to the reversal (in the same syringe) will help to circulate the drug to the patient.