

Weight (Kgs)	TTDEX VOLUME	MELOXICAM	ANTISEDAN (if necessary)	Weight (Lbs)
	mL (IM)	mL (SQ)	mL (IM)	
	0.035 mL/kg	0.1 mg/kg	1/3 Volume of TTDex	
	N/A	5 mg/mL	5 mg/mL	
1.0 - 1.1	0.04	0.02	0.01	2.2 - 2.4
1.2 - 1.3	0.04	0.03	0.01	2.6 - 2.9
1.4 - 1.5	0.05	0.03	0.02	3.1 - 3.3
1.6 - 1.7	0.06	0.03	0.02	3.5 - 3.7
1.8 - 2.0	0.07	0.04	0.02	4.0 - 4.4
2.1 - 2.2	0.08	0.04	0.03	4.6 - 4.8
2.3 - 2.4	0.08	0.05	0.03	5.1 - 5.3
2.5 - 2.7	0.09	0.05	0.03	5.5 - 5.9
2.8 - 2.9	0.10	0.06	0.03	6.2 - 6.4
3.0 - 3.1	0.11	0.06	0.04	6.6 - 6.8
3.2 - 3.4	0.12	0.07	0.04	7.0 - 7.5
3.5 - 3.6	0.12	0.07	0.04	7.7 - 7.9
3.7 - 3.8	0.13	0.08	0.04	8.1 - 8.4
3.9 - 4.0	0.14	0.08	0.05	8.6 - 8.8
4.1 - 4.3	0.15	0.08	0.05	9.0 - 9.5
4.4 - 4.5	0.16	0.09	0.05	9.7 - 9.9
4.6 - 4.7	0.16	0.09	0.05	10.1 - 10.3
4.8 - 5.0	0.17	0.10	0.06	10.6 - 11.0
5.1 - 5.2	0.18	0.10	0.06	11.2 - 11.4
5.3 - 5.4	0.19	0.11	0.06	11.7 - 11.9
5.5 - 5.6	0.19	0.11	0.06	12.1 - 12.3
5.7 - 5.9	0.20	0.12	0.07	12.5 - 13.0
6.0 - 6.1	0.21	0.12	0.07	13.2 - 13.4
6.2 - 6.4	0.22	0.13	0.07	13.6 - 14.1

Add 0.2 mL of saline to each dose of Antisedan for reversal
 Meloxicam 0.1 mg/kg administered SQ at prep
 TTDex dosed at 0.035 mL/kg

Torbagesic	10 mg/mL
Telazol (when reconstituted)	100 mg/mL
Dexdomitor	0.5 mg/mL

Purchase unconstituted Telazol vials, and reconstitute with 2.5 mL of Dexdomitor (500 mcg per mL) and 2.5 mL of Torbagesic (10 mg per mL). Give 0.035 mL/kg IM for total injectable anesthesia.

If feline reversal is necessary after greater than 60 minutes (patient not showing signs of recovery), Antisedan may be administered IM. The patient must be warm and hydrated, and also adding 0.2 mL of saline to the reversal (in same syringe) will help to circulate the drug to the patient.